

Geometric quantization of symplectic foliations

G.SARDANASHVILY

Department of Theoretical Physics, Physics Faculty, Moscow State University, 117234 Moscow, Russia, e-mail: sard@grav.phys.msu.su

Abstract.

Geometric quantization of a Poisson manifold need not imply quantization of its symplectic leaves. We provide the leafwise geometric quantization of a Poisson manifold, seen as a foliated one, whose quantum algebra restricted to each leaf is quantized.

Mathematics Subject Classification (2000): 53D50

Key words: leafwise differential calculus, leafwise cohomology, symplectic foliation, geometric quantization.

1 Introduction

Though there is one-to-one correspondence between the (regular) Poisson structures on a smooth manifold and its symplectic foliations, geometric quantization of a Poisson manifold need not imply quantization of its symplectic leaves [14].

Firstly, contravariant connections fail to admit the pull-back operation. Therefore, prequantization of a Poisson manifold does not determine straightforwardly prequantization of its symplectic leaves. Secondly, polarization of a Poisson manifold is defined in terms of sheaves of functions, and it need not be associated to any distribution. As a consequence, its pull-back onto a leaf is not polarization of a symplectic manifold in general. Thirdly, a quantum algebra of a Poisson manifold contains the centre of a Poisson algebra. However, there are models where quantization of this centre has no physical meaning. For instance, the centre of the Poisson algebra of a mechanical system with classical parameters consists of functions of these parameters.

Geometric quantization of symplectic foliations disposes of these problems. The quantum algebra of a symplectic foliation is also the quantum algebra of the associated Poisson manifold such that its restriction to each symplectic leaf is defined and quantized. Thus, geometric quantization of a symplectic foliation provides the leafwise quantization of a Poisson manifold. This is the case of systems whose symplectic leaves are indexed by non-quantizable variables, e.g., systems depending on classical parameters and constraint systems.

Geometric quantization of a symplectic foliation is phrased in terms of the leafwise differential calculus and leafwise connections on a foliated manifold. We show the following. Firstly, homomorphisms of the de Rham complex of a Poisson manifold both to its Lichnerowicz–Poisson complex and the de Rham complex of its symplectic leaf factorize

through the leafwise de Rham complex, and their cohomology groups do so. Secondly, any leafwise connection comes from a connection. Using these facts, we state the equivalence of prequantization of a Poisson manifold to prequantization of its symplectic foliation, which also yields prequantization of each symplectic leaf. On the contrary, polarization of a symplectic foliation is associated to a particular polarization of a Poisson manifold, and its restriction to any symplectic leaf is polarization of this leaf. Therefore, we define metaplectic correction of a symplectic foliation so that its quantum algebra restricted to each leaf is quantized. It is represented by Hermitian operators in the pre-Hilbert space of leafwise half-forms, integrable over the leaves of this foliation.

For example, the configuration space of a mechanical system with classical parameters is a fibre bundle $Q \rightarrow \Sigma$ over a manifold of parameters Σ [5, 6, 9]. Its momentum phase space is the vertical cotangent bundle V^*Q of $Q \rightarrow \Sigma$ endowed with the canonical Poisson structure, whose characteristic foliation coincides with the fibration $V^*Q \rightarrow \Sigma$. One can think of its fibre $V_\sigma^*Q = T^*Q_\sigma$, $\sigma \in \Sigma$, as being the momentum phase space of a mechanical system with fixed parameters. Of course, if a system is conservative, it can be quantized separately at each point of a parameter space, but the leafwise quantization procedure need be called into play if parameters are variable [9]

2 The leafwise differential calculus

Manifolds throughout are assumed to be smooth, Hausdorff, second-countable (i.e., paracompact), and connected.

A (regular) foliation $\mathcal{F}$ of a manifold Z consists of (maximal) integral manifolds of an involutive distribution $i_{\mathcal{F}} : T\mathcal{F} \rightarrow TZ$ on Z [8]. A foliated manifold $(Z, \mathcal{F})$ admits an adapted coordinate atlas

$$\{(U_\xi; z^\lambda; z^i)\}, \quad \lambda = 1, \dots, \text{codim } \mathcal{F}, \quad i = 1, \dots, \dim \mathcal{F}, \quad (1)$$

such that transition functions of coordinates z^λ are independent of the remaining coordinates z^i and, for each leaf F of a foliation $\mathcal{F}$, the connected components of $F \cap U_\xi$ are given by the equations $z^\lambda = \text{const}$. These connected components and coordinates (z^i) on them make up a coordinate atlas of a leaf F .

The real Lie algebra $\mathcal{T}_1(\mathcal{F})$ of global sections of the distribution $T\mathcal{F} \rightarrow Z$ is a $C^\infty(Z)$ -submodule of the derivation module of the $\mathbf{R}$ -ring $C^\infty(Z)$ of smooth real functions on Z . Its kernel $S_{\mathcal{F}}(Z) \subset C^\infty(Z)$ consists of functions constant on leaves of $\mathcal{F}$. Therefore, $\mathcal{T}_1(\mathcal{F})$ is the Lie $S_{\mathcal{F}}(Z)$ -algebra of derivations of $C^\infty(Z)$, regarded as a $S_{\mathcal{F}}(Z)$ -ring. Then one can introduce the leafwise differential calculus [4] as the Chevalley–Eilenberg differential calculus over the $S_{\mathcal{F}}(Z)$ -ring $C^\infty(Z)$. It is defined as a subcomplex

$$0 \rightarrow S_{\mathcal{F}}(Z) \longrightarrow C^\infty(Z) \xrightarrow{\tilde{d}} \mathfrak{F}^1(Z) \cdots \xrightarrow{\tilde{d}} \mathfrak{F}^{\dim \mathcal{F}}(Z) \rightarrow 0 \quad (2)$$

of the Chevalley–Eilenberg complex of the Lie $S_{\mathcal{F}}(Z)$ -algebra $\mathcal{T}_1(\mathcal{F})$ with coefficients in $C^\infty(Z)$ which consists of $C^\infty(Z)$ -multilinear skew-symmetric maps $\times^r \mathcal{T}_1(\mathcal{F}) \rightarrow C^\infty(Z)$, $r = 1, \dots, \dim \mathcal{F}$ [6]. These maps are global sections of exterior products $\wedge^r T\mathcal{F}^*$ of the dual $T\mathcal{F}^* \rightarrow Z$ of $T\mathcal{F} \rightarrow Z$. They are called the leafwise forms on a foliated manifold $(Z, \mathcal{F})$, and are given by the coordinate expression

$$\phi = \frac{1}{r!} \phi_{i_1 \dots i_r} \tilde{d}z^{i_1} \wedge \dots \wedge \tilde{d}z^{i_r},$$

where $\{\tilde{d}z^i\}$ are the duals of the holonomic fibre bases $\{\partial_i\}$ for $T\mathcal{F}$. Then one can think of the Chevalley–Eilenberg coboundary operator

$$\tilde{d}\phi = \tilde{d}z^k \wedge \partial_k \phi = \frac{1}{r!} \partial_k \phi_{i_1 \dots i_r} \tilde{d}z^k \wedge \tilde{d}z^{i_1} \wedge \dots \wedge \tilde{d}z^{i_r}$$

as being the leafwise exterior differential. Accordingly, (2) is called the leafwise de Rham complex (or the tangential de Rham complex in the terminology of [4]). This is the complex $(\mathcal{A}^{0,*}, d_f)$ in [11]. Its cohomology $H_{\mathcal{F}}^*(Z)$, called the leafwise de Rham cohomology, equals the cohomology $H^*(Z; S_{\mathcal{F}})$ of Z with coefficients in the sheaf $S_{\mathcal{F}}$ of germs of elements of $S_{\mathcal{F}}(Z)$ [2, 7].

Let us consider the exact sequence

$$0 \rightarrow \text{Ann } T\mathcal{F} \longrightarrow T^*Z \xrightarrow{i_{\mathcal{F}}^*} T\mathcal{F}^* \rightarrow 0. \quad (3)$$

of vector bundles over Z . Since it admits a splitting, the epimorphism $i_{\mathcal{F}}^*$ yields an epimorphism of the graded algebra $\mathcal{O}^*(Z)$ of exterior forms on Z to the algebra $\mathfrak{F}^*(Z)$ of leafwise forms. It obeys the condition $i_{\mathcal{F}}^* \circ d = \tilde{d} \circ i_{\mathcal{F}}^*$ and, thereby, provides the cochain morphism

$$i_{\mathcal{F}}^* : (\mathbf{R}, \mathcal{O}^*(Z), d) \rightarrow (S_{\mathcal{F}}(Z), \mathcal{F}^*(Z), \tilde{d}), \quad dz^\lambda \mapsto 0, \quad dz^i \mapsto \tilde{d}z^i, \quad (4)$$

of the de Rham complex of Z to the leafwise de Rham complex (2) and the corresponding homomorphism

$$[i_{\mathcal{F}}^*]^* : H^*(Z) \rightarrow H_{\mathcal{F}}^*(Z) \quad (5)$$

of the de Rham cohomology of Z to the leafwise one. Note that $[i_{\mathcal{F}}^*]^{r>0}$ need not be epimorphisms [11].

Given a leaf $i_F : F \rightarrow Z$ of a foliation $\mathcal{F}$, we have the pull-back homomorphism

$$(\mathbf{R}, \mathcal{O}^*(Z), d) \rightarrow (\mathbf{R}, \mathcal{O}^*(F), d) \quad (6)$$

of the de Rham complex of Z to that of F and the corresponding homomorphism of the de Rham cohomology groups

$$H^*(Z) \rightarrow H^*(F). \quad (7)$$

PROPOSITION 1. *The homomorphisms (6) – (7) factorize through the homomorphisms (4) – (5).*

Proof. It is readily observed that the pull-back bundles $i_F^*T\mathcal{F}$ and $i_F^*T\mathcal{F}^*$ over F are isomorphic to the tangent and the cotangent bundles of F , respectively. Moreover, a direct computation shows that $i_F^*(\tilde{d}\phi) = d(i_F^*\phi)$ for any leafwise form ϕ . It follows that the cochain morphism (6) factorizes through the cochain morphism (4) and the cochain morphism

$$i_F^* : (S_{\mathcal{F}}(Z), \mathcal{F}^*(Z), \tilde{d}) \rightarrow (\mathbf{R}, \mathcal{O}^*(F), d), \quad \tilde{d}z^i \mapsto dz^i, \quad (8)$$

of the leafwise de Rham complex of $(Z, \mathcal{F})$ to the de Rham complex of F . Accordingly, the cohomology morphism (7) factorizes through the leafwise cohomology

$$H^*(Z) \xrightarrow{[i_{\mathcal{F}}^*]} H_{\mathcal{F}}^*(Z) \xrightarrow{[i_F^*]} H^*(F). \quad (9)$$

□

Turn now to symplectic foliations. Let $\mathcal{F}$ be an even dimensional foliation of a manifold Z . A $\tilde{d}$ -closed non-degenerate leafwise two-form $\Omega_{\mathcal{F}}$ on a foliated manifold $(Z, \mathcal{F})$ is called symplectic. Its pull-back $i_F^*\Omega_{\mathcal{F}}$ onto each leaf F of $\mathcal{F}$ is a symplectic form on F . If a leafwise symplectic form $\Omega_{\mathcal{F}}$ exists, it yields the bundle isomorphism

$$\Omega_{\mathcal{F}}^b : T\mathcal{F} \xrightarrow{Z} T\mathcal{F}^*, \quad \Omega_{\mathcal{F}}^b : v \mapsto -v \lrcorner \Omega_{\mathcal{F}}(z), \quad v \in T_z\mathcal{F}. \quad (10)$$

The inverse isomorphism $\Omega_{\mathcal{F}}^\sharp$ determines the bivector field

$$w_{\Omega}(\alpha, \beta) = \Omega_{\mathcal{F}}(\Omega_{\mathcal{F}}^\sharp(i_{\mathcal{F}}^*\alpha), \Omega_{\mathcal{F}}^\sharp(i_{\mathcal{F}}^*\beta)), \quad \forall \alpha, \beta \in T_z^*Z, \quad z \in Z, \quad (11)$$

on Z subordinate to $\overset{2}{\wedge} T\mathcal{F}$. It is a Poisson bivector field (see the relation (20) below). The corresponding Poisson bracket reads

$$\{f, f'\}_{\mathcal{F}} = \vartheta_f \lrcorner \tilde{d}f', \quad \vartheta_f \lrcorner \Omega_{\mathcal{F}} = -\tilde{d}f, \quad \vartheta_f = \Omega_{\mathcal{F}}^\sharp(\tilde{d}f), \quad f, f' \in C^\infty(Z). \quad (12)$$

Its kernel is $S_{\mathcal{F}}(Z)$.

Conversely, let (Z, w) be a (regular) Poisson manifold and $\mathcal{F}$ its characteristic foliation. Since $\text{Ann } T\mathcal{F} \subset T^*Z$ is precisely the kernel of a Poisson bivector field w , the bundle homomorphism $w^\sharp : T^*Z \xrightarrow{Z} TZ$ factorizes in a unique fashion

$$w^\sharp : T^*Z \xrightarrow{Z} T\mathcal{F}^* \xrightarrow{Z} T\mathcal{F} \xrightarrow{Z} TZ \quad (13)$$

through the bundle isomorphism

$$w_{\mathcal{F}}^\sharp : T\mathcal{F}^* \xrightarrow{Z} T\mathcal{F}, \quad w_{\mathcal{F}}^\sharp : \alpha \mapsto -w(z) \lrcorner \alpha, \quad \alpha \in T_z\mathcal{F}^*. \quad (14)$$

The inverse isomorphism $w_{\mathcal{F}}^b$ yields the symplectic leafwise form

$$\Omega_{\mathcal{F}}(v, v') = w(w_{\mathcal{F}}^b(v), w_{\mathcal{F}}^b(v')), \quad \forall v, v' \in T_z \mathcal{F}, \quad z \in Z. \quad (15)$$

The formulae (11) and (15) establish the above mentioned equivalence between the Poisson structures on a manifold Z and its symplectic foliations, though this equivalence need not be preserved under morphisms.

Let us consider the Lichnerowicz–Poisson (henceforth LP) complex

$$0 \longrightarrow S_{\mathcal{F}}(Z) \longrightarrow C^{\infty}(Z) \xrightarrow{\widehat{w}} \mathcal{T}_1(Z) \xrightarrow{\widehat{w}} \cdots \xrightarrow{\widehat{w}} \mathcal{T}_{\dim \mathcal{F}}(Z) \rightarrow 0 \quad (16)$$

of multivector fields on a Poisson manifold (Z, w) with respect to the contravariant exterior differential

$$\widehat{w} : \mathcal{T}_r(Z) \rightarrow \mathcal{T}_{r+1}(Z), \quad \widehat{w}(\vartheta) = -[w, \vartheta], \quad \vartheta \in \mathcal{T}_*(Z),$$

where $[\cdot, \cdot]$ denotes the Schouten–Nijenhuis bracket. There are the cochain morphism

$$\begin{aligned} w^{\sharp} : (\mathbf{R}, \mathcal{O}^*(Z), d) &\rightarrow (S_{\mathcal{F}}(Z), \mathcal{T}_*(Z), -\widehat{w}) \\ w^{\sharp}(\phi)(\sigma_1, \dots, \sigma_r) &= (-1)^r \phi(w^{\sharp}(\sigma_1), \dots, w^{\sharp}(\sigma_r)), \quad \phi \in \mathcal{O}^r(Z), \quad \sigma_i \in \mathcal{O}^1(Z), \\ \widehat{w} \circ w^{\sharp} &= -w^{\sharp} \circ d. \end{aligned} \quad (17)$$

of the de Rham complex to the LP one and the corresponding homomorphism

$$[w^{\sharp}] : H^*(Z) \rightarrow H_w^*(Z) \quad (18)$$

of the de Rham cohomology of Z to the LP cohomology of the complex (15) [13].

PROPOSITION 2. *The cochain morphism $w^{\sharp}$ (17) factorizes through the leafwise complex (2) and, accordingly, the cohomology homomorphism $[w^{\sharp}]$ (18) does through the leafwise cohomology*

$$H^*(Z) \xrightarrow{[i_{\mathcal{F}}^*]} H_{\mathcal{F}}^*(Z) \longrightarrow H_w^*(Z). \quad (19)$$

Proof. Let $\mathcal{T}_*(\mathcal{F}) \subset \mathcal{T}_*(Z)$ denote the exterior subalgebra of multivector fields on Z subordinate to $T\mathcal{F}$, where $\mathcal{T}_0(\mathcal{F}) = C^{\infty}(Z)$. Clearly, $(S_{\mathcal{F}}(Z), \mathcal{T}_*(\mathcal{F}), \widehat{w})$ is a subcomplex of the LP complex (16). Since

$$\widehat{w} \circ \Omega_{\mathcal{F}}^{\sharp} = -\Omega_{\mathcal{F}}^{\sharp} \circ \widetilde{d}, \quad (20)$$

the bundle isomorphism $w_{\mathcal{F}}^{\sharp} = \Omega_{\mathcal{F}}^{\sharp}$ (14) yields the cochain isomorphism

$$\Omega_{\mathcal{F}}^{\sharp} : (S_{\mathcal{F}}(Z), \mathfrak{F}^*(Z), \widetilde{d}) \rightarrow (S_{\mathcal{F}}(Z), \mathcal{T}_*(\mathcal{F}), -\widehat{w})$$

of the leafwise de Rham complex (2) to the subcomplex $(\mathcal{T}_*(\mathcal{F}), \hat{w})$ of the LP complex (16). Then the composition

$$i_{\mathcal{F}} \circ \Omega_{\mathcal{F}}^{\sharp} : (S_{\mathcal{F}}(Z), \mathfrak{F}^*(Z), \tilde{d}) \rightarrow (S_{\mathcal{F}}(Z), \mathcal{T}_*(Z), -\hat{w}) \quad (21)$$

is a cochain monomorphism of the leafwise de Rham complex to the LP one (16). In view of the factorization (13), the cochain morphism (17) factorizes through the cochain morphisms (4) and (21). Accordingly, the cohomology homomorphism $[w^{\sharp}]$ (18) factorizes through the cohomology homomorphism $[i_{\mathcal{F}}^*]$ (5) and the cohomology homomorphism

$$[i_{\mathcal{F}} \circ \Omega_{\mathcal{F}}^{\sharp}] : H_{\mathcal{F}}^*(Z) \rightarrow H_w^*(Z). \quad (22)$$

□

3 Prequantization of a symplectic foliation

Prequantization of a symplectic foliation $(\mathcal{F}, \Omega_{\mathcal{F}})$ of a manifold Z provides a representation

$$f \mapsto i\hat{f}, \quad [\hat{f}, \hat{f}'] = -i\{\widehat{f}, \widehat{f'}\}_{\mathcal{F}}, \quad (23)$$

of the Poisson algebra $(C^{\infty}(Z), \{f, f'\}_{\mathcal{F}})$ by first order differential operators on sections of a complex line bundle $\pi : C \rightarrow Z$. These operators are given by the Kostant–Souriau formula

$$\hat{f} = -i\nabla_{\vartheta_f}^{\mathcal{F}} + \varepsilon f, \quad \vartheta_f = \Omega_{\mathcal{F}}^{\sharp}(\tilde{d}f), \quad \varepsilon > 0, \quad (24)$$

where $\nabla^{\mathcal{F}}$ is a leafwise connection on $C \rightarrow Z$ such that its curvature form obeys the prequantization condition

$$\tilde{R} = i\varepsilon\Omega_{\mathcal{F}}. \quad (25)$$

Using the fact that any leafwise connection comes from a connection (see Theorem 5 below), we will provide the cohomology analysis of this condition, and will show that prequantization of a symplectic foliation yields prequantization of its symplectic leaves.

DEFINITION 3. *In the framework of the leafwise differential calculus, a (linear) leafwise connection on the complex line bundle $C \rightarrow Z$ is defined as an algebraic connection $\nabla^{\mathcal{F}}$ on the $C^{\infty}(Z)$ -module $C(Z)$ of global sections of this bundle, where $C^{\infty}(Z)$ is regarded as a $S_{\mathcal{F}}(Z)$ -ring. It associates to each element $\tau \in \mathcal{T}_1(\mathcal{F})$ an $S_{\mathcal{F}}(Z)$ -linear endomorphism $\nabla_{\tau}^{\mathcal{F}}$ of $C(Z)$ which obeys the Leibniz rule*

$$\nabla_{\tau}^{\mathcal{F}}(fs) = (\tau \rfloor \tilde{d}f)s + f\nabla_{\tau}^{\mathcal{F}}(s), \quad f \in C^{\infty}(Z), \quad s \in C(Z). \quad (26)$$

Recall that a linear connection on $C \rightarrow Z$ can equivalently be defined as an algebraic connection on the module $C(Z)$ which assigns to each vector field $\tau \in \mathcal{T}_1(Z)$ on Z an $\mathbf{R}$ -linear endomorphism of $C(Z)$ obeying the Leibniz rule (26). Restricted to $\mathcal{T}_1(\mathcal{F})$, it obviously yields a leafwise connection. In order to show that any leafwise connection is of this form, we will appeal to an alternative definition of a leafwise connection in terms of leafwise forms.

The inverse images $\pi^{-1}(F)$ of leaves F of the foliation $\mathcal{F}$ of Z provide a (regular) foliation $C_{\mathcal{F}}$ of the line bundle C . Given the (holomorphic) tangent bundle $TC_{\mathcal{F}}$ of this foliation, we have the exact sequence of vector bundles

$$0 \rightarrow VC \xrightarrow{C} TC_{\mathcal{F}} \xrightarrow{C} C \times_Z T\mathcal{F} \rightarrow 0, \quad (27)$$

where VC is the (holomorphic) vertical tangent bundle of $C \rightarrow Z$.

DEFINITION 4. *A (linear) leafwise connection on the complex line bundle $C \rightarrow Z$ is a splitting of the exact sequence (27) which is linear over C .*

One can choose an adapted coordinate atlas $\{(U_{\xi}; z^{\lambda}; z^i)\}$ (1) of a foliated manifold $(Z, \mathcal{F})$ such that U_{ξ} are trivialization domains of the complex line bundle $C \rightarrow Z$. Let $(z^{\lambda}; z^i; c)$, $c \in \mathbf{C}$, be the corresponding bundle coordinates on $C \rightarrow Z$. They are also adapted coordinates on the foliated manifold $(C, C_{\mathcal{F}})$. With respect to these coordinates, a (linear) leafwise connection is represented by a $TC_{\mathcal{F}}$ -valued leafwise one-form

$$A_{\mathcal{F}} = \tilde{d}z^i \otimes (\partial_i + A_i c \partial_c), \quad (28)$$

where A_i are local complex functions on C .

The exact sequence (27) is obviously a subsequence of the exact sequence

$$0 \rightarrow VC \xrightarrow{C} TC \xrightarrow{C} C \times_Z TZ \rightarrow 0,$$

where TC is the holomorphic tangent bundle of C . Consequently, any connection

$$\Gamma = dz^{\lambda} \otimes (\partial_{\lambda} + \Gamma_{\lambda} c \partial_c) + dz^i \otimes (\partial_i + \Gamma_i c \partial_c) \quad (29)$$

on the complex line bundle $C \rightarrow Z$ yields a leafwise connection

$$\Gamma_{\mathcal{F}} = \tilde{d}z^i \otimes (\partial_i + \Gamma_i c \partial_c). \quad (30)$$

THEOREM 5. *Any leafwise connection on the complex line bundle $C \rightarrow Z$ comes from a connection on it.*

Proof. Let $A_{\mathcal{F}}$ (28) be a leafwise connection on $C \rightarrow Z$ and $\Gamma_{\mathcal{F}}$ (30) a leafwise connection which comes from some connection Γ (29) on $C \rightarrow Z$. Their affine difference over C is a section

$$Q = A_{\mathcal{F}} - \Gamma_{\mathcal{F}} = \tilde{d}z^i \otimes (A_i - \Gamma_i) c \partial_c$$

of the vector bundle $T\mathcal{F}^* \otimes_C VC \rightarrow C$. Given some splitting

$$B : \tilde{dz}^i \mapsto dz^i - B_\lambda^i dz^\lambda \quad (31)$$

of the exact sequence (3), the composition

$$(B \otimes \text{Id}_{VC}) \circ Q = (dz^i - B_\lambda^i dz^\lambda) \otimes (A_i - \Gamma_i) c \partial_c : C \rightarrow T^*Z \otimes_C VC$$

is a soldering form on the complex line bundle $C \rightarrow Z$. Then

$$\Gamma + (B \otimes \text{Id}_{VC}) \circ Q = dz^\lambda \otimes (\partial_\lambda + [\Gamma_\lambda - B_\lambda^i (A_i - \Gamma_i)] c \partial_c) + dz^i \otimes (\partial_i + A_i c \partial_c)$$

is a desired connection on $C \rightarrow Z$ which yields the leafwise connection $A_{\mathcal{F}}$ (28). $\square$

In particular, it follows that, in view of the above mentioned algebraic definition of a linear connection on a vector bundle, Definition 3 and Definition 4 of a leafwise connection are equivalent, namely,

$$\nabla^{\mathcal{F}} s = \tilde{d}s - A_i s \tilde{dz}^i, \quad s \in C(Z).$$

The curvature of a leafwise connection $\nabla^{\mathcal{F}}$ is defined as a $C^\infty(Z)$ -linear endomorphism

$$\tilde{R}(\tau, \tau') = \nabla_{[\tau, \tau']}^{\mathcal{F}} - [\nabla_\tau^{\mathcal{F}}, \nabla_{\tau'}^{\mathcal{F}}] = \tau^i \tau'^j R_{ij}, \quad R_{ij} = \partial_i A_j - \partial_j A_i, \quad (32)$$

of $C(Z)$ for any vector fields $\tau, \tau' \in \mathcal{T}_1(\mathcal{F})$. It is represented by the complex leafwise two-form

$$\tilde{R} = \frac{1}{2} R_{ij} \tilde{dz}^i \wedge \tilde{dz}^j. \quad (33)$$

If a leafwise connection $\nabla^{\mathcal{F}}$ comes from a connection ∇ , its curvature leafwise form $\tilde{R}$ (33) is the image $\tilde{R} = i_{\mathcal{F}}^* R$ of the curvature form R of the connection ∇ with respect to the morphism $i_{\mathcal{F}}^*$ (4).

Now let us turn to the prequantization condition (25).

LEMMA 6. *Let us assume that there exists a leafwise connection $\Gamma_{\mathcal{F}}$ on the complex line bundle $C \rightarrow Z$ which fulfils the prequantization condition (25). Then, for any Hermitian form g on $C \rightarrow Z$, there exists a leafwise connection $A_{\mathcal{F}}^g$ on $C \rightarrow Z$ which: (i) satisfies the condition (25), (ii) preserves g , and (iii) comes from a $U(1)$ -principal connection on $C \rightarrow Z$.*

Proof. For any Hermitian form g on $C \rightarrow Z$, there exists an associated bundle atlas $\Psi^g = \{(z^\lambda; z^i, c)\}$ of C with $U(1)$ -valued transition functions such that $g(c, c') = c\bar{c}'$. Let the above mentioned leafwise connection $\Gamma_{\mathcal{F}}$ comes from a linear connection Γ (29) on $C \rightarrow Z$ written with respect to the atlas Ψ^g . The connection Γ is split into the sum $A^g + \gamma$ where

$$A^g = dz^\lambda \otimes (\partial_\lambda + \text{Im}(\Gamma_\lambda) c \partial_c) + dz^i \otimes (\partial_i + \text{Im}(\Gamma_i) c \partial_c) \quad (34)$$

is a $U(1)$ -principal connection, preserving the Hermitian form g . The curvature forms R of Γ and R^g of A^g obey the relation $R^g = \text{Im}(R)$. The connection A^g (34) defines the leafwise connection

$$A_{\mathcal{F}}^g = i_{\mathcal{F}}^* A = \tilde{d}z^i \otimes (\partial_i + iA_i^g c \partial_c), \quad iA_i^g = \text{Im}(\Gamma_i), \quad (35)$$

preserving the Hermitian form g . Its curvature fulfils a desired relation

$$\tilde{R}^g = i_{\mathcal{F}}^* R^g = \text{Im}(i_{\mathcal{F}}^* R) = i\varepsilon \Omega_{\mathcal{F}}. \quad (36)$$

□

Since A^g (34) is a $U(1)$ -principal connection, its curvature form R^g is related to the first Chern form of integer de Rham cohomology class by the formula $c_1 = i(2\pi)^{-1} R^g$. If the prequantization condition (25) holds, the relation (36) shows that the leafwise cohomology class of the leafwise form $(2\pi)^{-1} \varepsilon \Omega_{\mathcal{F}}$ is the image of an integer de Rham cohomology class with respect to the cohomology morphism $[i_{\mathcal{F}}^*]$ (5). Conversely, if a leafwise symplectic form $\Omega_{\mathcal{F}}$ on a foliated manifold $(Z, \mathcal{F})$ is of this type, there exists a complex line bundle $C \rightarrow Z$ and a $U(1)$ -principal connection A on $C \rightarrow Z$ such that the leafwise connection $i_{\mathcal{F}}^* A$ fulfils the relation (25). Thus, we have stated the following.

PROPOSITION 7. *A symplectic foliation $(\mathcal{F}, \Omega_{\mathcal{F}})$ of a manifold Z admits prequantization (24) iff the leafwise cohomology class of $(2\pi)^{-1} \varepsilon \Omega_{\mathcal{F}}$ is the image of an integer de Rham cohomology class of Z .*

In particular, let (Z, w) be a Poisson manifold and $(\mathcal{F}, \Omega_{\mathcal{F}})$ its characteristic symplectic foliation. As is well-known, a Poisson manifold admits prequantization iff the LP cohomology class of the bivector field $(2\pi)^{-1} \varepsilon w$, $\varepsilon > 0$, is the image of an integer de Rham cohomology class with respect to the cohomology morphism $[w^{\sharp}]$ (18) [12, 13]. By virtue of Proposition 2, this morphism factorizes through the cohomology morphism $[i_{\mathcal{F}}^*]$ (5). Therefore, in accordance with Proposition 7, prequantization of a Poisson manifold takes place iff prequantization of its symplectic foliation does well, and both these prequantizations utilize the same prequantization bundle $C \rightarrow Z$. Herewith, each leafwise connection $\nabla^{\mathcal{F}}$ obeying the prequantization condition (25) yields the admissible contravariant connection $\nabla_{\phi}^w = \nabla_{w^{\sharp}(\phi)}^{\mathcal{F}}$, $\phi \in \mathcal{O}^1(Z)$, on $C \rightarrow Z$ whose curvature bivector equals $i\varepsilon w$. Clearly, $\nabla^{\mathcal{F}}$ and ∇^w lead to the same prequantization formula (24).

Let F be a leaf of a symplectic foliation $(\mathcal{F}, \Omega_{\mathcal{F}})$ provided with the symplectic form $\Omega_F = i_F^* \Omega_{\mathcal{F}}$. In accordance with Proposition 1 and the commutative diagram of cohomology groups

$$\begin{array}{ccc} H^*(Z; \mathbf{Z}) & \longrightarrow & H^*(Z) \\ \downarrow & & \downarrow \\ H^*(F; \mathbf{Z}) & \longrightarrow & H^*(F) \end{array},$$

the symplectic form $(2\pi)^{-1}\varepsilon\Omega_F$ belongs to an integer de Rham cohomology class if a leafwise symplectic form $\Omega_{\mathcal{F}}$ fulfils the condition of Proposition 7. This states the following.

PROPOSITION 8. *If a symplectic foliation admits prequantization, each its symplectic leaf does well.*

The corresponding prequantization bundle for F is the pull-back complex line bundle i_F^*C , coordinated by (z^i, c) . Furthermore, let $A_{\mathcal{F}}^g$ (35) be a leafwise connection on the prequantization bundle $C \rightarrow Z$ which obeys Lemma 6, i.e., comes from a $U(1)$ -principal connection A^g on $C \rightarrow Z$. Then the pull-back

$$A_F = i_F^*A^g = dz^i \otimes (\partial_i + ii_F^*(A_i^g)c\partial_c) \quad (37)$$

of the connection A^g onto $i_F^*C \rightarrow F$ satisfies the prequantization condition

$$R_F = i_F^*R = i\varepsilon\Omega_F,$$

and preserves the pull-back Hermitian form i_F^*g on $i_F^*C \rightarrow F$.

4 Polarization of a symplectic foliation

Let us define polarization of a symplectic foliation $(\mathcal{F}, \Omega_{\mathcal{F}})$ of a manifold Z as a maximal (regular) involutive distribution $\mathbf{T} \subset T\mathcal{F}$ on Z such that

$$\Omega_{\mathcal{F}}(u, v) = 0, \quad \forall u, v \in \mathbf{T}_z, \quad z \in Z. \quad (38)$$

Given the Lie algebra $\mathbf{T}(Z)$ of $\mathbf{T}$ -subordinate vector fields on Z , let $\mathcal{A}_{\mathcal{F}} \subset C^\infty(Z)$ be the complexified subalgebra of functions f whose leafwise Hamiltonian vector fields ϑ_f (12) fulfil the condition $[\vartheta_f, \mathbf{T}(Z)] \subset \mathbf{T}(Z)$. It is called the quantum algebra of a symplectic foliation $(\mathcal{F}, \Omega_{\mathcal{F}})$ with respect to the polarization $\mathbf{T}$. This algebra obviously contains the centre $S_{\mathcal{F}}(Z)$ of the Poisson algebra $(C^\infty(Z), \{, \}_{\mathcal{F}})$, and is a Lie $S_{\mathcal{F}}(Z)$ -algebra.

PROPOSITION 9. *Every polarization $\mathbf{T}$ of a symplectic foliation $(\mathcal{F}, \Omega_{\mathcal{F}})$ yields polarization of the associated Poisson manifold (Z, w_Ω) .*

Proof. Let us consider the presheaf of local smooth functions f on Z whose leafwise Hamiltonian vector fields ϑ_f (12) are subordinate to $\mathbf{T}$. The sheaf Φ of germs of these functions is polarization of the Poisson manifold (Z, w_Ω) . Equivalently, Φ is the sheaf of germs of functions on Z whose leafwise differentials are subordinate to the codistribution $\Omega_{\mathcal{F}}^\perp \mathbf{T}$. $\square$

Note that the polarization Φ need not be maximal, unless $\mathbf{T}$ is of maximal dimension $\dim \mathcal{F}/2$. It belongs to the following particular type of polarizations of a Poisson manifold. Since the cochain morphism $i_{\mathcal{F}}^*$ (4) is an epimorphism, the leafwise differential calculus $\mathfrak{F}^*$ is universal, i.e., the leafwise differentials $\tilde{d}f$ of functions $f \in C^\infty(Z)$ on Z make up a basis for

the $C^\infty(Z)$ -module $\mathfrak{F}^1(Z)$. Let $\Phi(Z)$ denote the structure $\mathbf{R}$ -module of global sections of the sheaf Φ . Then the leafwise differentials of elements of $\Phi(Z)$ make up a basis for the $C^\infty(Z)$ -module of global sections of the codistribution $\Omega_{\mathcal{F}}^b \mathbf{T}$. Equivalently, the leafwise Hamiltonian vector fields of elements of $\Phi(Z)$ constitute a basis for the $C^\infty(Z)$ -module $\mathbf{T}(Z)$. Then one can easily show that polarization $\mathbf{T}$ of a symplectic foliation $(\mathcal{F}, \Omega_{\mathcal{F}})$ and the corresponding polarization Φ of the Poisson manifold (Z, w_Ω) in Proposition 9 define the same quantum algebra $\mathcal{A}_{\mathcal{F}}$.

Let (F, Ω_F) be a symplectic leaf of a symplectic foliation $(\mathcal{F}, \Omega_{\mathcal{F}})$. Given a polarization $\mathbf{T} \rightarrow Z$ of $(\mathcal{F}, \Omega_{\mathcal{F}})$, its restriction $\mathbf{T}_F = i_F^* \mathbf{T} \subset i_F^* T\mathcal{F} = TF$ to F is an involutive distribution on F . It obeys the condition

$$i_F^* \Omega_{\mathcal{F}}(u, v) = 0, \quad \forall u, v \in \mathbf{T}_{Fz}, \quad z \in F,$$

i.e., is polarization of the symplectic manifold (F, Ω_F) . Thus, we have stated the following.

PROPOSITION 10. *Polarization of a symplectic foliation defines polarization of each symplectic leaf.*

Clearly, the quantum algebra $\mathcal{A}_F$ of a symplectic leaf F with respect to the polarization $\mathbf{T}_F$ contains all elements $i_F^* f$ of the quantum algebra $\mathcal{A}_{\mathcal{F}}$ restricted to F .

5 Quantization of a symplectic foliation

Since $\mathcal{A}_{\mathcal{F}}$ is the quantum algebra both of a symplectic foliation $(\mathcal{F}, \Omega_{\mathcal{F}})$ and the associated Poisson manifold (Z, w_Ω) , one let us start from the standard metaplectic correction technique [1, 15].

Assuming that Z is oriented and that $H^2(Z; \mathbf{Z}_2) = 0$, let us consider the metilinear complex line bundle $\mathcal{D} \rightarrow Z$ characterized by an atlas $\Psi_Z = \{(U_\xi; z^\lambda; z^i; c)\}$ with the transition functions $c' = Sc$ such that $S\overline{S}$ is the inverse Jacobian of coordinate transition functions on Z . Global sections of this bundle are half-forms on Z . The metilinear bundle $\mathcal{D}$ belongs to the category of natural bundles, and the Lie derivative

$$\mathbf{L}_\tau = \tau^\lambda \partial_\lambda + \tau^i \partial_i + \frac{1}{2}(\partial_\lambda \tau^\lambda + \partial_i \tau^i) \quad (39)$$

of its sections along any vector field τ on Z is defined. The quantization bundle is the tensor product $Y = C \otimes \mathcal{D}$. The space $Y_K(Z)$ of its sections of compact support is provided with the non-degenerate Hermitian form

$$\langle \rho | \rho' \rangle = \left(\frac{1}{2\pi} \right)^{\dim Z/2} \int_Z \rho \rho', \quad \rho, \rho' \in Y_K(Z), \quad (40)$$

written with respect to the atlases Ψ^g of C and Ψ_Z of $\mathcal{D}$. Given the leafwise connection $A_{\mathcal{F}}^g$ (35) and the Lie derivative $\mathbf{L}$ (39), one can assign the first order differential operator

$$\hat{f} = -i[(\nabla_{\vartheta_f}^{\mathcal{F}} + i\varepsilon f) \otimes \text{Id} + \text{Id} \otimes \mathbf{L}_{\vartheta_f}] = -i[\nabla_{\vartheta_f}^{\mathcal{F}} + i\varepsilon f + \frac{1}{2}\partial_i \vartheta_f^i], \quad f \in \mathcal{A}_{\mathcal{F}}, \quad (41)$$

on $Y_K(Z)$ to each element of the quantum algebra $\mathcal{A}_{\mathcal{F}}$. These operators obey the Dirac condition (23), and provide a representation of the quantum algebra $\mathcal{A}_{\mathcal{F}}$ by (unbounded) Hermitian operators in the pre-Hilbert space $Y_K(Z)$. Finally, this representation is restricted to the subspace E of sections $\rho \in Y_K(Z)$ which obey the condition

$$(\nabla_{\vartheta}^{\mathcal{F}} \otimes \text{Id} + \text{Id} \otimes \mathbf{L}_{\vartheta})\rho = (\nabla_{\vartheta}^{\mathcal{F}} + \frac{1}{2}\partial_i \vartheta^i)\rho = 0$$

for all $\mathbf{T}$ -subordinate leafwise Hamiltonian vector fields ϑ .

However, it may happen that the above quantization has no physical sense because the Hermitian form (40) on the carrier space E and, consequently, the mean values of operators (41) are defined by integration over the whole manifold Z . For instance, it implies integration over classical parameters. Therefore, we suggest a different scheme of quantization of symplectic foliations.

Let us consider the exterior bundle $\wedge^m T\mathcal{F}^*$, $m = \dim \mathcal{F}$. Its structure group $GL(m, \mathbf{R})$ is reducible to the group $GL^+(m, \mathbf{R})$ since a symplectic foliation is oriented. One can regard this fibre bundle as being associated to a $GL(m, \mathbf{C})$ -principal bundle $P \rightarrow Z$. As earlier, let us assume that $H^2(Z; \mathbf{Z}_2) = 0$. Then the principal bundle P admits a two-fold covering principal bundle with the structure metlinear group $ML(m, \mathbf{C})$ [1]. As a consequence, there exists a complex line bundle $\mathcal{D}_{\mathcal{F}} \rightarrow Z$ characterized by an atlas $\Psi_{\mathcal{F}} = \{(U_{\xi}; z^{\lambda}; z^i; c)\}$ with the transition functions $c' = S_{\mathcal{F}}c$ such that

$$S_{\mathcal{F}}\overline{S}_{\mathcal{F}} = \det \left(\frac{\partial z^i}{\partial z'^j} \right).$$

One can think of its sections as being leafwise half-forms on Z . The metlinear bundle $\mathcal{D}_{\mathcal{F}} \rightarrow Z$ admits the canonical lift of any $\mathbf{T}$ -subordinate vector field τ on Z . The corresponding Lie derivative of its sections reads

$$\mathbf{L}_{\tau}^{\mathcal{F}} = \tau^i \partial_i + \frac{1}{2} \partial_i \tau^i. \quad (42)$$

We define the quantization bundle as the tensor product $Y_{\mathcal{F}} = C \otimes \mathcal{D}_{\mathcal{F}}$. Given a leafwise connection $A_{\mathcal{F}}^g$ (35) and the Lie derivative $\mathbf{L}^{\mathcal{F}}$ (42), let us associate the first order differential operator

$$\hat{f} = -i[(\nabla_{\vartheta_f}^{\mathcal{F}} + i\varepsilon f) \otimes \text{Id} + \text{Id} \otimes \mathbf{L}_{\vartheta_f}^{\mathcal{F}}] = -i[\nabla_{\vartheta_f}^{\mathcal{F}} + i\varepsilon f + \frac{1}{2}\partial_i \vartheta_f^i], \quad f \in \mathcal{A}_{\mathcal{F}}, \quad (43)$$

on sections $\rho_{\mathcal{F}}$ of $Y_{\mathcal{F}}$ to each element of the quantum algebra $\mathcal{A}_{\mathcal{F}}$. A direct computation with respect to the local Darboux coordinates on Z proves the following.

LEMMA 11. *The operators (43) obey the Dirac condition (23).*

LEMMA 12. *If a section $\rho_{\mathcal{F}}$ fulfils the condition*

$$(\nabla_{\vartheta}^{\mathcal{F}} \otimes \text{Id} + \text{Id} \otimes \mathbf{L}_{\vartheta}^{\mathcal{F}})\rho_{\mathcal{F}} = (\nabla_{\vartheta}^{\mathcal{F}} + \frac{1}{2}\partial_i\vartheta^i)\rho_{\mathcal{F}} = 0 \quad (44)$$

for all $\mathbf{T}$ -subordinate leafwise Hamiltonian vector field ϑ , then $\widehat{f}\rho_{\mathcal{F}}$ for any $f \in \mathcal{A}_{\mathcal{F}}$ possesses the same property.

Let us restrict the representation of the quantum algebra $\mathcal{A}_{\mathcal{F}}$ by the operators (43) to the subspace $E_{\mathcal{F}} \in Y_{\mathcal{F}}(Z)$ of sections $\rho_{\mathcal{F}}$ which obey the condition (44) and whose restriction to any leaf of $\mathcal{F}$ is of compact support. The last condition is motivated by the following.

Since $i_F^*T\mathcal{F}^* = T^*F$, the pull-back $i_F^*\mathcal{D}_{\mathcal{F}}$ of $\mathcal{D}_{\mathcal{F}}$ onto a leaf F is a metilinear bundle of half-forms on F . By virtue of Proposition 8 and Proposition 10, the pull-back $i_F^*Y_{\mathcal{F}}$ of the quantization bundle $Y_{\mathcal{F}} \rightarrow Z$ onto F is a quantization bundle for the symplectic manifold $(F, i_F^*\Omega_{\mathcal{F}})$. Given the pull-back connection A_F (37) and the polarization $\mathbf{T}_F = i_F^*\mathbf{T}$, this symplectic manifold is subject to the standard geometric quantization by the first order differential operators

$$\widehat{f} = -i(i_F^*\nabla_{\vartheta_f}^{\mathcal{F}} + i\varepsilon f + \frac{1}{2}\partial_i\vartheta_f^i), \quad f \in \mathcal{A}_{\mathcal{F}}, \quad (45)$$

on sections ρ_F of $i_F^*Y_{\mathcal{F}} \rightarrow F$ of compact support which obey the condition

$$(i_F^*\nabla_{\vartheta}^{\mathcal{F}} + \frac{1}{2}\partial_i\vartheta^i)\rho_F = 0 \quad (46)$$

for all $\mathbf{T}_F$ -subordinate Hamiltonian vector fields ϑ on F . These sections constitute a pre-Hilbert space E_F with respect to the Hermitian form

$$\langle \rho_F | \rho'_F \rangle = \left(\frac{1}{2\pi} \right)^{m/2} \int_F \rho_F \rho'_F.$$

The key point is the following.

PROPOSITION 13. *We have $i_F^*E_{\mathcal{F}} \subset E_F$, and the relation*

$$i_F^*(\widehat{f}\rho_{\mathcal{F}}) = (\widehat{i_F^*f})(i_F^*\rho_{\mathcal{F}}) \quad (47)$$

holds for all elements $f \in \mathcal{A}_{\mathcal{F}}$ and $\rho_{\mathcal{F}} \in E_{\mathcal{F}}$.

Proof. One can use the fact that the expressions (45) and (46) have the same coordinate form as the expressions (43) and (44) where $z^\lambda = \text{const}$. $\square$

The relation (47) enables one to think of the operators $\hat{f}$ (43) in $E_{\mathcal{F}}$ as being the leafwise quantization of the $S_{\mathcal{F}}(Z)$ -algebra $\mathcal{A}_{\mathcal{F}}$ in the pre-Hilbert $S_{\mathcal{F}}(Z)$ -module.

For example, the instantwise quantization of time-dependent mechanics is of this type [3, 10].

References

- [1] Echeverría Enríquez, A., Muñoz Lecanda, M., Román Roy, N. and Victoria-Monge, C.: *Extracta Math.* **13** (1998), 135.
- [2] El Kasimi-Alaoui, A.: *Compositio Math.* **49** (1983), 195.
- [3] Giachetta, G., Mangiarotti, L. and Sardanashvily G.: *J. Math. Phys.* **42** (2001) No.12.
- [4] Hector, G., Macías, E. and Saralegi, M.: *Publ. Mat.* **33** (1989), 423.
- [5] Mangiarotti, L. and Sardanashvily, G.: *Gauge Mechanics*, World Scientific, Singapore, 1998.
- [6] Mangiarotti, L. and Sardanashvily, G.: *Connections in Classical and Quantum Field Theory*, World Scientific, Singapore, 2000.
- [7] Mostov, M.: *Mem. Amer. Math. Soc.* **7** (1976), No.175.
- [8] Reinhart, B.: *Differential Geometry and Foliations*, Springer-Verlag, Berlin, 1983.
- [9] Sardanashvily, G.: *J. Math. Phys.* **41** (2000), 5245.
- [10] Śniatycki, J.: *Geometric Quantization and Quantum Mechanics*, Springer-Verlag, Berlin, 1980.
- [11] Vaisman, I.: *Cohomology and Differential Forms*, Marcel Dekker, Inc., New York, 1973.
- [12] Vaisman, I.: *J. Math. Phys.* **32** (1991), 3339.
- [13] Vaisman, I.: *Lectures on the Geometry of Poisson Manifolds*, Birkhäuser Verlag, Basel, 1994.
- [14] Vaisman, I.: *Diff. Geom. Appl.* **7** (1997), 265.
- [15] Woodhouse, N.: *Geometric Quantization*, Clarendon Press, Oxford, 1992.